

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

X-621-70-77
PREPRINT

NASA TM X-63905

DISCREPANCIES IN THE OBSERVED "PLASMA-TROUGH" DENSITY

E. J. R. MAIER
G. P. SERBU

MARCH 1970



GSFC

GODDARD SPACE FLIGHT CENTER
GREENBELT, MARYLAND

N 70-27302

FACILITY FORM 602	(ACCESSION NUMBER)	(THRU)
	14	1
	(PAGES)	(CODE)
	TMX 63905	25
	(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)

DISCREPANCIES IN THE OBSERVED "PLASMA-TROUGH" DENSITY

by

E.J.R. Maier and G.P. Serbu
Laboratory for Planetary Atmospheres
Goddard Space Flight Center
Greenbelt, Maryland

ABSTRACT

Observations of the plasmopause have now been achieved by a variety of techniques - ground based reception of Whistlers, and in-situ observations by mass spectrometers, retarding analyzers, and VLF receivers. Comparisons of some of these observations are made and it is concluded that the details of the plasmasphere boundary and the thermal plasma density beyond the plasmasphere are more variable than the idealized plasmasphere model would suggest.

Carpenter, in a classic paper published in 1966, described the plasma surrounding the earth as contained in a "plasmasphere" having a sharply defined outer boundary, beyond which the plasma density was of the order of $1/\text{cm}^3$. His results were derived from Whistler dispersion observations which are somewhat dependent on the model plasma distribution assumed within the plasmasphere and which tend to emphasize the plasma distribution near the magnetic equatorial plane. In addition, the acquisition of Whistler data from a specific region in space is dependent upon the probability of propagation of Whistler through the particular region.

Recently a number of spacecraft carrying retarding analyzers, mass spectrometers, and VLF receivers have traversed the plasmasphere/plasma trough region obtaining data along specified

trajectory paths. In this note we compare results obtained by various experimenters which demonstrate that the plasma density in the trough region may not always be the 1 ion/cm^3 stated in the idealized model.

In a recent paper Harris, Sharp and Chappell [1970] have criticized the IMP II electron results [Serbu and Maier, 1966a] for ... "not [having] always clearly shown the presence of the plasmopause". Their criticism is along the same general lines expressed by Gringauz [1969]... "In the interpretation of the data from this experiment (IMP II electron measurement) the authors (op. cit.) did not detect the knee in the distribution of charged particle density". These criticisms are based in part on our statement ... "as can be seen, neither of the density profiles of Figure 4 and 5 indicate a knee-type of fall-off in the region from 3 to 5 R_E ; in fact, our density of 5 R_E is just about a factor of 10 above that reported by Angerami and Carpenter [1966]".

Experimental evidence obtained first with the Pioneer I Quadrispherical Plasma Probe by Wolfe et al. [1967] and later with the Explorer 35 Retarding Potential Analyzer by Serbu [1969] indicates that as a consequence of solar particle and ultraviolet radiation a secondary electron sheath may envelop a sunlit spacecraft. The density of the photoelectrons can exceed the density of the ambient electrons and as a consequence, a valid measurement is not easily made in regions of very low electron density. Whipple and Parker [1969a] have developed

the theoretical response function for a Retarding Potential Analyzer experiment operating in a tenuous plasma in sunlight. They show that the response characteristic for the electron current-voltage (I-V) curve contains a primary, or ambient, component and a secondary emission component. Using experimental data from the OGO-I spacecraft, in the solar wind region, they also show that some of the secondary electrons are in ballistic trajectories escaping from and returning to the spacecraft, and moving with thermal energies in the range from 1 to 5 eV. Correcting our IMP-II [op. cit.] electron density values for the secondary component will not significantly change the shape of the electron density profiles within the plasmasphere; however, it will result in density values lower than originally reported for the region beyond the plasmopause. Structure in the density at the plasmopause is evident in data published by us [Serbu and Maier, 1966a], but due to the editorial requirement for brevity these profiles were omitted from the Journal of Geophysical Research publication. Note that Carpenter's [1966] original definition for the "knee" was a variation in the plasma density of 100 to 1 over a radial distance increment of $0.1 R_E$. We used this definition in noting the absence of a "knee" in our Figure 4 and 5 [op. cit.] data.

There is a large body of published data which we cite in support of our claim [Serbu and Maier, 1970] that the plasma density on the far side of the plasmopause exceeds the value

of 1 electron/cm³ given by Carpenter [op. cit.], see Figure 1. Chappell et al. [1970] claim to establish as a final value a density of <10 cm⁻³ for the plasma trough region. In Figure 2 we present their spectrometer data from OGO-5 which clearly shows a density of 100 ions/cm³ at L = 9.

Carpenter et al. [1969] presented the data of Figure 3 which shows the simultaneous V.L.F. and ion mass spectrometer data used to detect the plasmopause on July 5, 1966. They have identified the plasmopause at 4.6 R_E, where the observed ion density drop-off is a factor of 8 over a radial distance of 0.5 R_E. The plasma trough density measured on July 5, 1966 was in the range from 100 to 20 ions/cm³, which is also at variance with the criteria set up by Chappell et al. [op. cit.] for the plasma trough value of <10 ions/cm³. In Figure 4 we present the density data of Bezrukikh [1968] obtained from Electron 4. In the author's own words ... "The distribution of n_i, presented in Figure 6, gives grounds to believe that at the geomagnetic latitudes $\phi > 45^\circ$ the actual distribution of charged particle density in some cases may considerably differ from the idealized model, suggested by Carpenter [1966]".

In Figure 5 we present data of Vasyliunas [1968] which we offer in heuristic support to our contention for the intermittent existence of higher densities in the trough region. The mean electron energy versus density data of OGO-I and Vela [Bame et al. 1967] indicate that within the region of the plasma trough the density at 10 eV should be in the range 15 to 100 cm⁻³. We see

no reason for assuming that the generalized Maxwellian distribution function derived by Vasyliunas to adequately describe the energy density relationship as measured from 10,000 eV to 30 eV should require an abrupt modification at 10 eV.

Thus, we conclude there is evidence to indicate that the morphology of the plasma trough is both time and space dependent. In addition the plasma trough position and the character of the boundary may be dependent upon the energy interval used in the observation. At equatorial latitudes thermal particles, viz. the plasma component observable through Whistlers, have a temporal and spatial structure which at times agrees with and at other times is markedly different from that observed at higher energies [Schield and Frank, 1969] or at greater latitudes, Bezrukikh op. cit. For particles of energies less than 10 eV, there may exist external mechanisms such as photoemissions, secondary emission and spacecraft potential changes which can bring about a shift in the energy interval of the observation. Whipple and Parker [1970] calculate that for experiments which sample the ambient plasma more than one order of magnitude effects can be caused by small changes in the spacecraft potential. Results from long term observations, Serbu and Maier op. cit. and Taylor [1970] indicate that a high degree of variability exists in the plasma trough structure, that is, in the density, density gradients and location of the drop-off.

LIST OF FIGURES

- Figure 1 - The idealized model for the Thermal Plasma ($E \leq 1.0$ eV) particle density within the Magnetosphere. In this presentation "Plasmapause" was used synonymously with "knee" by Carpenter [1966].
- Figure 2 - The Magnetosphere ion density as a function of L, observed with a light ion mass spectrometer aboard the OGO-5 spacecraft. Chappell et al. [1970].
- Figure 3 - The simultaneous V.L.F. and ion mass spectrometer data which were used to detect the plasmapause on July 5, 1966. Carpenter et al. [1969].
- Figure 4 - The Electron 4 measurements of ion density as a function of radial distance; these observations were conducted at geomagnetic latitudes $\varphi > 45^\circ$. Bezrukikh [1968].
- Figure 5 - The density of Low-Energy Electrons ($10 \text{ KeV} \leq E \leq 30 \text{ eV}$) in the "Plasma Sheet" region of the Magnetosphere observed with the OGO-I and Vela satellites. Lines of constant energy density for three values of 'equivalent field strength' are also shown. Vasyliunas [1968].

REFERENCES

- Angerami, J.J. and D.L. Carpenter, Whistler Studies of the Plasmopause in the Magnetosphere -2., J. Geophys. Res., 71, 111-125, 1966.
- Bame, J.R., H. E. Asbridge, H.E. Felthausen, E.W. Hones, and I.B. Strong, Characteristics of the Plasma Sheet in the Earth's Magnetotail, J. Geophys. Res., Vol. 72, 113-129, 1967.
- Bezrukikh, V.V., Results of Measurements of the Density of Charged Particles in the Earth's Plasma Envelope Carried Out From Satellites Electron 2 and Electron 4, presented at the International Symposium on the Physics of the Magnetosphere, Washington, D.C., Sept. 3-13, 1968. Submitted to Kosmich. Issled., 1968.
- Carpenter, D.L., Whistler Studies of the Plasmopause in the Magnetosphere -1., J. Geophys. Res., 71, 693-709, 1966.
- Carpenter, D.L., and C.G. Park, H.A. Taylor, Jr., and H.C. Brinton, Multi-Experiment Detection of the Plasmopause from EOGO Satellites and Antarctic Ground Stations, J. Geophys. Res., 74, 1837-1847, 1969.
- Chappell, C.R., K.K. Harris, and G.W. Sharp, A Study of the Influence of Magnetic Activity on the Location of the Plasmopause as Measured by OGO 5, J. Geophys. Res., 75, 50-56, 1970.
- Gringauz, K.I., Low-Energy Plasma in the Earth's Magnetosphere, Magnetospheric Physics, edited by Donald J. Williams and Gilbert D. Mead. Published by the American Geophysical Union as Reviews of Geophysics, Vol. 7,

February-May, 1969.

Harris, K.K., G.W. Sharp, and C.R. Chappell, Observations of the Plasmapause from OGO 5, J. Geophys. Res., 75, 219-224, 1970.

Serbu, G.P. and E.J.R. Maier, Low-Energy Electrons Measured on IMP 2, J. Geophys. Res., 71, 3755-3766, 1966a.

Serbu, G.P., E.J.R. Maier, Low Energy Electrons Measured on IMP-II, Goddard Space Flight Center document X615-66-92, January, 1966b.

Serbu, G.P., Explorer 35 Measurements of Low-Energy Plasma in Lunar Orbit, J. Geophys. Res., 74, 372-376, 1969.

Serbu, G.P. and E.J.R. Maier, Observations from OGO-V of the Thermal Ion Density and Temperature within the Magnetosphere, submitted J. Geophys. Res., also published as a Goddard Space Flight Center Document, X615-70-34, 1970.

Schild, M.A., and L.A. Frank, Inner Edge of the Plasma Sheet, Submitted to J. Geophys. Res., University of Iowa Document No. 69-46.

Taylor, H.A., private communications, 1970.

Vasyliunas, V. M., A Survey of Low-Energy Electrons in the Evening Sector of the Magnetosphere with OGO 1 and OGO 3, J. Geophys. Res., 73, 2839-2884, 1968.

Wolfe, J. H., R. W. Silva, and D.D. McKibbin, The Interplanetary Solar Wind Electron Characteristics, Trans. Am. Geophys. Union, 48, 182, 1967.

Whipple, E.C. Jr., and L.W. Parker, Theory of an Electron Trap on a Charged Spacecraft, J. Geophys. Res., 74, 2962-2971, 1969.

**Whipple, E.C. Jr., and L.W. Parker, Effects of Secondary Electron
Emission on Electron Trap Measurements in the Magnetosphere
and Solar Wind, J. Geophys. Res., 74, 5763-5774, 1969.**




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5